

# HAMILTON, BROOK, SMITH & REYNOLDS, P.C.

## UTILITY PATENT APPLICATION TRANSMITTAL

*(Only for new nonprovisional applications under  
37 C.F.R. 1.53(b))*

Attorney Docket No.

YFL98-01pA

First Named Inventor or  
Application Identifier

Ya Fang Liu

Express Mail Label No.

EL136701351US

Title of  
Invention

METHOD FOR IDENTIFYING JNK AND MLK INHIBITORS FOR TREATMENT  
OF NEUROLOGICAL CONDITIONS

### APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents  
Box Patent Application  
Washington, D.C. 20231

1. ☐ Fee Transmittal Form  
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages  ]  
(preferred arrangement set forth below)
  - Descriptive title of the invention
  - Cross References to Related Applications
  - Statement Regarding Fed sponsored R & D
  - Reference to microfiche Appendix
  - Background of the Invention
  - Summary of the Invention
  - Brief Description of the Drawings
  - Detailed Description
  - Claim(s)
  - Abstract of the Disclosure
3. ☒ Drawing(s) (35 U.S.C. 113) [Total Sheets  ]
4. ☐ Oath or Declaration/POA [Total Pages  ]
  - a. ☐ Newly executed (original or copy)
  - b. ☐ Copy from a prior application (37 C.F.R. 1.63(d))  
(for continuation/divisional with Box 17 completed)  
[NOTE Box 5 below]
    - i. ☐ DELETION OF INVENTOR(S)  
Signed statement attached deleting  
inventor(s) named in the prior application,  
see 37 C.F.R. 1.63(d)(2) and 1.33(b).
5. ☐ Incorporation By Reference (useable if Box 4b is checked)  
The entire disclosure of the prior application, from which a copy  
of the oath or declaration is supplied under Box 4b, is considered  
as being part of the disclosure of the accompanying application  
and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. ☐ Nucleotide and/or Amino Acid Sequence Submission  
(if applicable, all necessary)
  - a. ☐ Computer Readable Copy
  - b. ☐ Paper Copy (identical to computer copy)
  - c. ☐ Statement verifying identity of above copies

### ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & documents)
9. ☐ 37 C.F.R. 3.73(b) Statement ☐ Power of Attorney  
(when there is an assignee)
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503) (2)  
(Should be specifically itemized)
14. ☐ Small Entity Statement(s) ☐ Statement filed in prior application,  
Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)
16. ☐ Other:.....

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: \_\_\_\_\_/\_\_\_\_\_

Prior application information: Examiner: \_\_\_\_\_ Group Art Unit: \_\_\_\_\_

### 18. CORRESPONDENCE ADDRESS

NAME	Alice O. Carroll, Esq.				
	HAMILTON, BROOK, SMITH & REYNOLDS, P.C.				
ADDRESS	Two Militia Drive				
CITY	Lexington	STATE	MA	ZIP CODE	02421-4799
COUNTRY	USA	TELEPHONE	(781) 861-6240	FAX	(781) 861-9540

Signature	<i>Alice O. Carroll</i>	Date	<i>September 17, 1998</i>
Submitted by Typed or Printed Name	Alice O. Carroll	Reg. Number	33,542

-1-

Date: 9/17/98 Express Mail Label No. EL136701351US

Inventor: Ya Fang Liu

Attorney's Docket No.: YFL98-01pA

METHOD FOR IDENTIFYING JNK AND MLK  
INHIBITORS FOR TREATMENT OF NEUROLOGICAL CONDITIONS

RELATED APPLICATION

5 This application claims the benefit of U.S. Provisional Application Serial No. 60/085,439, filed May 14, 1998, the entire teachings of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

10 Excitotoxicity is related to excessive activation of glutamate receptors which results in neuronal cell death. The physiological function of glutamate receptors is the mediation of ligand-gated cation channels with the concomitant influx of calcium, sodium and potassium through this receptor-gated channel. The influx of these cations is essential for maintaining membrane potentials and the plasticity of neurons which in  
15 itself plays a pivotal role in cognitive function of the central nervous system. Li, H.B., *et al.*, *Behav. Brain Res.*, 83:225-228 (1997); Roesler, R., *et al.*, *Neurology*, 50:1195 (1998); Wheal, H.V., *et al.*, *Prog. Neurobiol.*, 55:611-640 (1998); Wangen, K., *et al.*, *Brain Res.*, 99:126-130 (1997). Excitotoxicity plays an important role in neuronal cell death following acute insults such as hypoxia, ischemia, stroke and trauma, and it also  
20 plays a significant role in neuronal loss in AIDS dementia, epilepsy, focal ischemia.

Coyle, J.T. & Puttfarcken, P., *Science*, 262:689-695 (1993). Neurodegenerative disorders, such as Huntington's disease (HD), Alzheimer's disease (AD), Parkinson's disease (PD) and amyotrophic lateral sclerosis (ALS), are characterized by the progressive loss of a specific population of neurons in the central nervous system.

- 5 Growing evidence suggests that glutamate-mediated excitotoxicity may be a common pathway which contributes to neuronal cell death in a wide range of neurological disorders. Coyle, J.T. & Puttfarcken, P., *Science*, 262:689-695 (1993).

The molecular mechanisms of excitotoxicity-mediated neuronal cell death remains obscure. Over-production of free radicals that lead to impairment of  
10 mitochondrial function is the most widely held hypothesis. Beal, M.F., *et al.*, *Ann. Neurol.*, 38:357-366 (1995); Coyle, J.T. & Puttfarcken, P., *Science*, 262:689-695 (1993). However, it is unclear whether the increase of free radicals is the precursor that initiates neuronal degeneration or, rather, a subsequent consequence of neuronal degeneration. Interestingly, administration of antioxidants has little neuroprotective effect in patients  
15 suffering from various neurodegenerative diseases. Shults, C.W., *et al.*, *Neurology*, 50:793-795 (1998). Thus, some other mechanism(s) must exist for excitotoxicity-induced neuronal cell death.

c-Jun N-terminal kinases (JNKs) are identified as kinases which are activated upon stimulation by various environmental stimuli such as UV light,  $\gamma$  irradiation and  
20 mitogenic signals. Hibi, M., *et al.*, *Genes Dev.*, 7:2135-2148 (1993); Kyriakis, J.M., *et al.*, *Nature*, 369:156-160 (1994). The precise biological function of JNKs remains to be explored. However, some recent reports suggest that JNKs are involved in neuronal apoptosis induced by deprivation of survival factors, i.e., neurotrophic factors which support neuronal survival. Ham, J., *et al.*, *Neuron*, 14:927-939 (1995).

25 Mixed-lineage kinases (MLKs), so called because these proteins contain structural domains associated with a variety of cell types, were cloned from a cDNA library derived from mRNA from cancer tissue. MLKs were initially thought to participate in the oncogenesis of some cancers, although high levels of expression of MLKs were

found in the normal brain. Dorow, D.S., *et al.*, *Eur. J. Biochem.*, 213:701-710 (1993); Dorow, D.S., *et al.*, *Eur. J. Biochem.*, 234:492-500 (1995).

Searching for biochemical targets which are amenable to screening for neuroprotective therapeutic agents is of central concern in neuroscience today.

5 However, no clinically available pharmaceutical tool to date is employed for blocking excitotoxicity and preventing neuronal cell loss in various neurological disorders due to a lack of suitable biochemical targets. Glutamate receptor antagonists, such as MK-801, although successful in protecting neurons in animal experiments, have all failed in the clinical setting due to their blockage of cognitive function mediated by the receptors, as  
10 well as high toxicity to the central nervous system. Thus, an understanding of the molecular mechanism(s) of neuronal cell death induced by excitotoxicity is essential for the identification of new biochemical targets and the establishment of reliable methods for screening new therapeutic drugs from chemical libraries that can be utilized in the treatment of a variety of neurological disorders.

15 SUMMARY OF THE INVENTION

This invention relates to the discovery that inhibiting a JNK or MLK within a hippocampal neuronal cell can protect the cell from apoptosis. As such, JNK and MLK can be used as drug targets to screen for therapeutic agents to prevent glutamate or  
20 kainic acid mediated toxicity, to block excitotoxicity and to prevent neuronal loss in a variety of neurological conditions, such as Huntington's disease and Alzheimer's disease.

In one aspect of the invention, a method is described for assessing a compound's ability to inhibit neuronal cell death, and thus to identify compounds that can be used to  
25 prevent and/or treat neurological conditions. According to the method, neuronal cells having activated MLK and/or JNK activity are contacted with a compound and the number of neuronal cells that die is determined. A decrease in the number of dead neuronal cells in the presence of the compound compared to the number of dead

neuronal cells in the absence of the compound is indicative of the compound's ability to inhibit neuronal cell death. Preferably, the neuronal cells are apoptotic neurons (i.e., cell death caused by a neurological condition) or neurons that are induced to undergo apoptosis, such as by contacting the neuronal cells with neurotoxin (e.g., glutamate, quinolinic acid or kainic acid); or by genetic manipulation of the neuronal cells. Most preferred are HN33 hippocampal neuronal cells.

In another embodiment, the invention features a method for testing a compound's potential as a drug for treating a mammal (e.g., a human) susceptible to or having a neurological condition by (1) contacting a compound with a JNK (e.g., JNK3) or MLK (e.g., MLK2); (2) measuring the level of a JNK-associated or MLK-associated activity (e.g., a kinase activity); and (3) comparing the level of the JNK-associated or MLK-associated activity in the presence of the compound with the level of the JNK-associated or MLK-associated activity in the absence of the compound. The compound is a potentially useful drug for treating the mammal when the level of the JNK-associated or MLK-associated activity in the presence of the compound is less than the level of the JNK-associated or MLK-associated activity in the absence of the compound.

The JNK or MLK can be within a cell, which can be an animal (e.g., human) cell *in vivo*. When the JNK or MLK is within a cell, the JNK-associated or MLK-associated activity can be apoptosis, which can be measured by a TUNEL assay (described below). Apoptosis within such a cell can be induced by introducing into the cell a huntingtin protein that has at least 40 consecutive glutamic acids (e.g., polyglutamine stretch-expanded huntingtin). Alternatively, apoptosis can be induced by introducing into the cell the C-terminal 100 amino acids of an amyloid precursor protein (APP). Preferably, the huntingtin protein or the amyloid precursor protein is introduced by a vector, especially a nucleic acid vector. When the cell is within an animal, the JNK-associated or MLK-associated activity can be neurodegeneration.

The invention also features a method for testing a compound's potential as a drug for treating a mammal (e.g., a human) susceptible to or having a neurological condition

by (1) contacting a compound with a neuronal cell containing a JNK (e.g., JNK3) or MLK (e.g., MLK2); (2) measuring the level of a JNK or MLK protein activity (e.g., kinase activity, such as the presence or amount of phosphorylated product) in the cell; and (3) comparing the level of the JNK or MLK protein activity in the cell in the presence of the compound with the level of the JNK or MLK protein activity in the cell in the absence of the compound. The compound is a potentially useful drug for treating the mammal when the level of the JNK or MLK protein activity in the cell in the presence of the compound is less than the level of the JNK or MLK protein activity in the cell in the absence of the compound. Alternatively, cell viability can be ascertained by determining the degree of neuronal cell death, wherein a decreased number of dead neuronal cells in the presence of the compound compared to the number of dead neuronal cells in the absence of the compound is indicative of the compound's ability to inhibit JNK or MLK protein activity, thereby preventing neuronal cell death.

In another aspect, the invention provides a method for testing the potential of a JNK or MLK inhibitor as a drug for treating a mammal (e.g., a human) susceptible to or having a neurological condition. The method can be performed on compounds identified as JNK and/or MLK inhibitory agents using the methods of this invention to confirm their inhibitory effectiveness under apoptotic conditions. Accordingly, the method provides (1) incubating a neuronal cell in the presence of a JNK or MLK inhibitor; (2) contacting surviving cells with an agent that induces apoptosis in the cell; and (3) comparing the occurrence of apoptosis in the cell in the presence of the JNK or MLK inhibitor with the occurrence of apoptosis in the cell in the absence of the JNK or MLK inhibitor. The compound is a potentially useful drug for treating the mammal when the occurrence of apoptosis in the cell in the presence of the JNK or MLK inhibitor is less than the occurrence of apoptosis in the cell in the absence of the JNK or MLK inhibitor.

The methods of the invention are used to identify inhibitors of JNK or MLK which are potentially useful for the treatment of a neurological condition, including

neuronal cell death following acute insults such as hypoxia, ischemia, stroke, and trauma. Other neurological conditions treatable with compounds identified by the methods of the invention include AIDS dementia, epilepsy, focal ischemia, Huntington's disease, Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. Each of these conditions are characterized by the progressive loss of a specific population of neurons in the central nervous system. The methods of the invention are particularly useful in finding compounds which can be used to prevent and/or treat neurological conditions, including genetic neurological conditions. The invention also pertains to compounds, identified using the methods described herein, that inhibit MLK and/or JNK activity and that prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, particularly neurodegenerative diseases, such as Huntington's disease and Alzheimer's disease.

The invention also provides methods for preventing and/or treating neuronal conditions in a mammal comprising administering to a mammal, in need thereof, an effective therapeutic amount of a compound that inhibits JNK and/or MLK. The inhibitory effects of the compound will reduce and/or prevent neuron cell death occurring in a mammal susceptible to or having a neurological condition. In a preferred embodiment, the neurological condition is a neurological disease whereby glutamate or kainic acid mediated excitotoxicity is involved in neuronal cell death. JNK and/or MLK inhibitors identified using any of the methods described herein are useful as therapeutic or prophylactic drugs to prevent neuronal loss.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph illustrating a time course of cell death upon expression of normal or mutated huntingtin in HN33 cells. pcDNA1 (solid box), pFL16HD (circles), pFL48HD (triangles) and pFL89HG (open boxes).

Fig. 2 is a graph illustrating apoptotic cell death induced by expression of mutated huntingtin with 48 or 89 polyglutamine repeats was completely blocked by added the

ICE or CPP32 inhibitor in the medium indicating HN33 cells are undergoing apoptotic cell death.

Fig. 3 is a graph illustrating the fold of increased JNK activity in HN33 cells upon expression of mutated huntingtin with 48 or 89 polyglutamine repeats.

5 Fig. 4 is a graph illustrating apoptotic cell death of HN33 cells induced by mutated huntingtin with either 48 or 89 polyglutamine repeats was inhibited by co-expression of dominant negative mutant form of SEK1 but not wild-type SEK1.

Fig. 5 is a graph illustrating apoptotic cell death of HN33 cells induced by the treatment of glutamate (250  $\mu$ M) or kainic acid (kainate, 200  $\mu$ M) was significantly  
10 attenuated by expression of dominant negative mutant form of SEK1 (K54R) but not wild-type SEK1, indicating that glutamate or kainate induced the activation of the SEK1-JNK pathway to mediate neuronal apoptosis.

Fig. 6 is a graph illustrating the fold of increased JNK activity in HN33 cells upon stimulation of glutamate or kainate receptor indicating that glutamate or kainate receptor  
15 activation also stimulated the JNK activity like expression of mutated huntingtin.

Fig. 7 is a graph illustrating a time course of cell death upon following transfection of pcDNA1 (control) APP or APP deletion mutant APP-C-100, pcDNA1 (open box), wild-type APP (solid diamond), APP-C-100 (solid circles).

Fig. 8 is a graph illustrating the fold of increased JNK activity in HN33 cells  
20 following transfection of pcDNA1 (control) APP or APP deletion mutant APP-C-100, indicating that expression of APP-C-100 stimulated the JNK activity.

Fig. 9 is a graph illustrating apoptotic cell death of HN33 cells induced by expression of APP-C-100 was significantly attenuated by co-expression of dominant negative mutant form of SEK1 (K54R) but not wild-type SEK1, indicating that amyloid  
25 precursor protein induced the activation of the SEK1-JNK pathway to mediate neuronal apoptosis.



Fig. 10 is a graph illustrating a time course of cell death following transfection of pRK5CMV (control), wild-type MLK2 or kinase dead version of MLK2, pRK5CMV (open box), wild-type MLK2 (solid diamond), kinase dead MLK2 (solid circles).

Fig. 11 is a graph illustrating apoptotic cell death of HN33 cells induced by expression of MLK2 was significantly attenuated by co-expression of dominant negative mutant form of SEK1 (K54R) but not wild-type SEK1, indicating that MLK2 induced the activation of the SEK1-JNK pathway to mediate neuronal apoptosis.

Fig. 12 is a graph illustrating apoptotic cell death of HN33 cells induced by expression mutated huntingtin with 48 or 89 CAG repeats was blocked by co-expression of kinase dead MLK2, indicating that the MLK2-associated activity mediated neuronal cell death in Huntington's diseases.

Fig. 13 is a graph illustrating apoptotic cell death of HN33 cells induced by the treatment of glutamate (250  $\mu$ M) or kainic acid (kainate 200  $\mu$ M) was blocked by expression of kinase dead MLK2 indicating that the MLK2-associated activity mediated neuronal cell death in neuronal excitotoxicity induced by glutamate or kainate receptor activation.

Fig. 14 is a graph illustrating apoptotic cell death of HN33 cells induced by expression of deletion APP mutant APP-C-100 was blocked by co-expression of kinase dead MLK2, indicating that the MLK2-associated activity mediated neuronal cell death in Alzheimer's diseases.

#### DETAILED DESCRIPTION OF THE INVENTION

The invention relates to the discovery that two families of proteins, JNK and MLK, can serve as targets for the treatment of neurological conditions. It has been discovered that inhibition of JNK3, a member of the JNK family, and MLK2, a member of the MLK family, can protect a neuronal cell from apoptosis induced by polyglutamine-expanded huntingtin, whose expression caused HD in humans. The Huntington's Disease Collaborative Research Group, *Cell*, 72:971-983 (1993). Thus,

the invention provides a method for assessing compounds for their potential as drugs for the treatment of neurological conditions, such as Huntington's disease, by determining whether the compound can inhibit a JNK- or MLK-associated activity.

5 MLK and JNK participate in a biochemical cascade (activation of MLK-SEK1-JNK1) that mediates neuronal cell toxicity. Upon stimulation by glutamate, kainic acid, or other receptor agonists, the glutamate receptor, located at the cell surface, is activated and interacts with associated proteins (e.g., PDZ domain-containing proteins) whose SH3 domain in turn binds to a MLK protein, thereby activating its kinase activity. The MLK protein directly binds to and stimulates a SEK1 protein which in turn binds to and stimulates JNK. Over stimulation of JNK can lead to neuronal apoptosis (cell death).  
10 Normally MLK is inhibited by a protein, such as normal huntingtin, binding to its SH3 domain thus inhibiting the enzyme's kinase activity. This inhibition prevents the formation of the cascade resulting in no or little JNK activity, thereby preserving neuronal cell viability.

15 To determine whether polyglutamine-expanded huntingtin induced cell toxicity in neurons, the expression of huntingtins with 16, 48 or 89 CAG repeats in an immortalized rat hippocampal neuronal cell line (HN33) was examined. HN33 has been described in detail in Hammond, D. N., *et al.*, *Brain Res.*, 512:190-200 (1990); the entire teachings of which are incorporated herein by reference. The hippocampal neurons serve as a potential target of mutated huntingtin, mutated amyloid precursor  
20 protein (APP), as well as glutamate stimulation. The results demonstrated that expression of mutated huntingtin with 48 or 89 CAG repeats stimulated JNKs and induced apoptotic cell death in HN33 cells, while expression of normal huntingtin with 16 CAG repeats had no toxic effect. JNK activation occurs several hours prior to neuronal apoptosis suggesting that it is an early signal for the induction of neuronal  
25 apoptosis. Furthermore, co-expression of a dominant negative form of stress signaling kinase (SEK1), which serve as a specific inhibitor of JNKs, attenuated both JNK activation and neuronal apoptosis induced by mutated huntingtin. This study

demonstrates that JNK activation can mediate neuronal cell death in neurological disorders. Like mutated huntingtin, the treatment of HN33 cells with glutamate also resulted in JNK activation and apoptotic cell death. Expression of dominant negative mutant SEK1 significantly attenuated glutamate-induced toxicity. Taken together, these studies indicate that JNK is a common cellular mediator for induction of neuronal cell death mediated by both glutamate and mutated huntingtin.

The mechanism for how mutated huntingtin stimulates JNKs was also examined. The huntingtin protein contains multiple SH3 domain binding sites. In previous studies, it has been demonstrated that normal huntingtin directly binds to SH3 domain-containing proteins Grb2 and RasGAP. Liu, Y.F., *et al.*, *J. Biol. Chem.*, 272:8121-8124 (1997). MLK1, MLK2 and MLK3 (MLKs) are the only known kinases that directly activate the SEK1-JNK cascade and contain a SH3 domain as well as a SH3 domain binding site. MLK2 is a neuronal form of MLKs. The presence of MLKs, in particular MLK2, appears to be required for mutated huntingtin-mediated JNK activation and neuronal apoptosis. MLKs were expressed in HN33 cells where expression of mutated huntingtin induced JNK activation and apoptosis, while in both 293 and CHO cells where MLKs are absent, the expression of mutated huntingtin did not generate any cell toxicity in 293 and CHO cells. However, co-expression of mutated huntingtin along with MLK2 in 293 cells caused rapid apoptotic cell death. The SH3 domain of MLKs is required for their proper cellular localization and activation of the SEK1-JNK pathway. In the *in vitro* binding studies, it was shown that normal huntingtin binds to the SH3 domain of MLK2 and inhibits enzyme activity while expansion of the polyglutamine repeats interferes with huntingtin's binding and consequently is not inhibitory with respect to MLK2's activity. Furthermore, expression of kinase dead MLK2 completely blocks mutated huntingtin-induced neuronal cell death. These studies indicate that mutation of huntingtin results in an increase of free MLKs thereby causing over-activation of the MLK-SEK1-JNK cascade which ultimately leads to neuronal cell death.

The SH3 binding motif is found in six other proteins involved in polyglutamine repeat-expanded neurodegenerative hereditary diseases, such as ataxia-1, ataxia-2, ataxia-6, ataxia-7, Kennedy disease, dentatorubral and pallidoluysian atrophy (DRPLA). The normal (wild-type) counter-part proteins bind to and suppress MLK activity, in  
 5 contrast to the mutated protein form which lose such ability, resulting in the over-activation of the MLK2-SEK1-JNK pathway in neurons.

The JNK kinases phosphorylate and activate the transcription factor c-Jun which mediates apoptosis. Recent studies suggest that c-Jun may serve as an important mediator for neuronal apoptosis induced by a variety of environmental stresses. In  
 10 primary cultured sympathetic, hippocampal or cerebellar granule neurons, deprivation of growth factor in these primary cultures lead to persistent activation of JNKs and consequently the phosphorylation of c-Jun. Ham, J., *et al.*, *Neuron*, 14:927-939 (1995); Xia, Z., *et al.*, *Science*, 270:1326-1330 (1995). Suppression of c-Jun expression by antisense-oligonucleotides, or functional blockade by microinjection of antibodies or  
 15 expression of dominant negative c-Jun prevents neuronal apoptosis; in contrast, over-expression of c-Jun induces apoptosis in sympathetic neurons. Estus, S., *et al.*, *J. Cell Biol.*, 127:1717-1727 (1994); Ham, J., *et al.*, *Neuron*, 14:927-939 (1995). Enhanced c-Jun expression occurs in degenerating and apoptotic neurons after ischemia, nerve fiber transection, axotomized brains, UV irradiation and various other types of  
 20 neuronal injury. Ferrer, I., *et al.*, *Eur. J. Neurosci.*, 8:1286-1298 (1998); Herdegen, T., *et al.*, *J. Neurosci.*, 18:5124-5135 (1998).

Increased c-Jun expression and activation are also implicated in the generation of the neuronal apoptotic process induced by glutamate or kainic acid. Administration of glutamate or quinolinic acid, a N-methyl-D-aspartate (NMDA) receptor agonist, or  
 25 kainic acid in rats results in a rapid induction of c-Jun expression and neuronal apoptosis. Coyle, J.T. & Puttfarcken, P., *Science*, 262:689-695 (1993); Qin, Z.-H., *et al.*, *Mol. Pharmacol.*, 53:33-42 (1997). These physiological events can be blocked by the

NMDA receptor antagonist MK-801. Qin, Z.-H., *et al.*, *Mol. Pharmacol.*, 53:33-42 (1997).

Over-activation of the c-Jun-mediated JNK cascade has also been implicated in various other neurodegenerative disorders. Enhanced c-Jun expression is observed in the brains from patients suffering from multiple sclerosis, AD and ALS. Anderson, A.J., *et al.*, *Exp. Neurol.*, 125:286-295 (1994); Martin, D.G., *et al.*, *Neurosci. Lett.*, 212:95-98 (1996). These studies support the basis of the present invention in that the over-activation (or stimulation) of the MLK-SEK1-JNK cascade leads to the increase of expression, activation and translocation of c-Jun which is responsible for neuronal cell death in these, and other, neurodegenerative disorders. Thus, inhibition of this cascade can protect neurons from toxicity induced by endo- and/or exo-toxins including, but not limited to, mutated proteins like huntingtin, quinolinic acid, kainic acid, glutamate over-excitation as well as other etiological agents.

Based upon these findings, JNK and MLK can be used as targets for the development of inhibitory compounds of JNK- and MLK-associated activity, and such compounds can be used to prevent neuronal loss, such as induced by excitotoxicity or glutamate- or kainic acid-mediated toxicity. As used herein, a “JNK-associated activity” is any biochemical, cellular, or physiological property that varies with any variation in JNK gene transcription or translation, or JNK protein activity. Likewise, a “MLK-associated activity” is any biochemical, cellular, or physiological property that varies with any variation in MLK gene transcription or translation, or MLK protein activity. A JNK or MLK inhibitor is a compound that inhibits a JNK or MLK protein activity. A JNK or MLK protein activity is any measurable biochemical activity possessed by the protein, e.g., a kinase activity or an ability to bind to another protein such as c-Jun.

Inhibitors of MLKs identified by the methods described herein can block persistent activation of glutamate receptor-induced over-activation of MLKs without affecting other receptor functions, such as the involvement of neuronal plasticity and

cognitive functions. Inhibition of MLKs will attenuate the JNK activity in neurons and protect neurons from excitotoxicity thereby preventing neuronal loss in these diseases. Inhibitors of JNKs or MLKs identified by the methods described herein can suppress glutamate receptor-induced activation of the MLK-SEK1-JNK cascade and prevent neuronal apoptosis in various neurological diseases.

The term "neurological condition" as used herein is intended to embrace disorders, disease states and disturbances which cause or result in neuronal cell injury, compromise or cell death. Neurological conditions can result from axonal degeneration, ischemia due to stroke, heart arrest, exposure, exposure to neurotoxins such as, but not limited to, glutamate, kainic acid and quinolinic acid, MPTP exposure to bacterial or viral toxins, impaired function or dysfunction of neurons such as increase or decrease of neurotransmitter synthesis and/or release. Neurological diseases and disturbances include, but are not limited to, Alzheimer's disease; Parkinson's disease; motor neuron diseases such as amyotrophic lateral sclerosis (ALS), Huntington's disease and syringomyelia; ataxias, dementias; chorea; dystonia; dysllesia; encephalomyelopathy; parenchymatous cerebellar degeneration; Kennedy disease; Down syndrome; progressive supranuclear palsy; DRPLA, stroke or other ischemic injuries; thoracic outlet syndrome, trauma; electrical brain injuries; decompression brain injuries; AIDS dementia; multiple sclerosis; epilepsy; concussive or penetrating injuries of the brain or spinal cord; peripheral neuropathy; brain injuries due to exposure of military hazards such as blast over-pressure, ionizing radiation, and genetic neurological conditions. By "genetic neurological condition" is meant a neurological condition, or a predisposition to it, that is caused at least in part by or correlated with a specific gene or mutation within that gene; for example, a genetic neurological condition can be caused by or correlated with more than one specific gene. Examples of genetic neurological conditions include, but are not limited to, Alzheimer's disease, Huntington's disease, spinal and bulbar muscular atrophy, fragile X syndrome, FRAXE mental retardation,

myotonic dystrophy, spinocerebellar ataxia type 1, dentatorubral-pallidoluysian atrophy, and Machado-Joseph disease.

In one aspect of the invention, a method is described for assessing a compound's ability to inhibit neuronal cell death. According to the method, neuronal cells having  
5 activated MLK and/or JNK activity are contacted with a compound and the number of neuronal cells that die is determined. A decrease in the number of dead neuronal cells in the presence of the compound compared to the number of dead neuronal cells in the absence of the compound is indicative of the compound's ability to inhibit neuronal cell death. Preferably, the neuronal cells are apoptotic neurons or neurons that are induced  
10 to undergo apoptosis neuronal cells with neurotoxin or genetic manipulation.

A neuronal cell useful in the methods of the invention is preferably susceptible to JNK-dependent or MLK-dependent apoptosis. To facilitate apoptosis, such a cell can express a polypeptide known to be associated with or induce a neurodegenerative disease, such as a polyglutamine-expanded polypeptide (e.g., polyglutamine-expanded  
15 huntingtin) or the C-terminal 100 amino acid fragment of an amyloid precursor protein. A preferred neuronal cell that is useful for assessing MLK and/or JNK inhibitors is an immortalized rat hippocampal neuronal cell line HN33, with or without genetic manipulations to induce apoptosis, as described above.

In another embodiment, the invention features a method for testing a compound's  
20 potential as a drug for treating a mammal (e.g., a human) susceptible to or having a neurological condition by (1) contacting a compound with a JNK (e.g., JNK3) or MLK (e.g., MLK2); (2) measuring the level of a JNK-associated or MLK-associated activity (e.g., a kinase activity); and (3) comparing the level of the JNK-associated or MLK-associated activity in the presence of the compound with the level of the JNK-associated  
25 or MLK-associated activity in the absence of the compound. The compound is a potentially useful drug for treating the mammal when the level of the JNK-associated or MLK-associated activity in the presence of the compound is less than the level of the JNK-associated or MLK-associated activity in the absence of the compound.

In another aspect of the invention, a putative inhibitory agent is incubated *in vitro* in the presence of JNK and appropriate JNK substrates, such as c-Jun and a phosphate donor like adenosine triphosphate (ATP), under conditions sufficient for enzymatic activity; followed by isolating the phosphorylated product. Isolated JNK protein, including JNK1, JNK2 and JNK3, can be obtained for this, as well as other assays, by several different molecular and chromatographic methods known to those skilled in the art. The JNK polypeptides useful in the methods of the present invention are preferably wild-type whose sequence is known and readily available. The human JNK3 polypeptide is described by Martin *et al.*, *Mol. Brain Res.*, 35:47-57 (1996). Other JNK proteins useful in the methods of the invention include those described in GenBank Accession Nos. U17743, U49249 and AF006689. Isolated JNK protein, from about 0.5 µg to about 2 µg of purified JNK, is incubated with substrate in an aqueous medium, such as a kinase buffer (containing about: 20 mM HEPES, pH 7.5, 15 mM MgCl<sub>2</sub>, 15 mM β-glycerophosphate, 0.1 mM Na<sub>2</sub>PO<sub>4</sub> and 2 mM dithiothreitol) at about 30°C for approximately 15 minutes. The substrates that can be used in this reaction include, but are not limited to, c-Jun, from about 1 µg to about 3 µg, a known substrate for JNK's kinase activity, and the phosphate donor, ATP (approximately 2.5 mM). For detection purposes, 5 µCi of [γ-<sup>32</sup>]ATP can be used as a co-substrate. The assay system can also include in the incubation mixture a putative inhibitory JNK agent. The reaction can be terminated by addition of Laemmli buffer, approximately 20 µL. The addition of this buffer will also prepare the sample for product analysis. The reaction mixture can be subjected to sodium dodecylsulfate polyacrylamide gel electrophoresis (hereinafter SDS-PAGE) in order to determine the amount of phosphorylated c-Jun that was formed in the reaction. The radioactivity emitted from the γ<sup>32</sup>P can be measured using conventional radioactivity gel detection systems, such as an X-ray followed by β-scan. The phosphorylated c-Jun product will have a different migration rate along the gel when compared to the labeled ATP co-substrate and therefore will not be confused with the kinase product. A determination can then be made concerning whether the test



agent inhibited JNK's activity by comparing reaction mixtures having the agent present to reaction mixtures without addition of the compound.

Alternatively, JNK substrates, such as c-Jun and ATP, can be incubated in the presence of a cellular extract containing putative JNK enzyme activity, including JNK1, JNK2 and JNK3. An inhibitory agent to be tested can be placed in the reaction vial along with the other reactants to examine the efficacy of the agent. The reaction and detection protocol can be conducted in the same manner as that describe above for the *in vitro* assay without cellular extract. The cellular extract can originate from a cell or tissue culture system, or can be prepared from whole tissue employing isolation and purification protocols known to those skilled in the art.

In another embodiment, the invention pertains to contacting a cell with a putative inhibitory agent in order to screen for inhibitory agents of JNK activity, including JNK1, JNK2 and JNK3. The cell to be contacted can be of a cell or tissue culture system. The putative inhibitory agent is delivered to the cell under conditions sufficient for enzymatic activity in any of a number of ways known to those skilled in the art. If the agent is not membrane permeable, then the agent can be delivered into the cell via electroporation, or if it is a polypeptide, a nucleic acid or viral vector can be employed. If the cell has JNK activity present in an active form, then JNK can be stimulated by delivering to the cell SEK1, a known stimulator of JNK. If the cell lacks a JNK gene or functional JNK gene or transcript or translational product, the cell can be transfected with an operatively linked JNK gene. "Operatively linked" is intended to mean that the nucleotide sequence is linked to a regulatory sequence in a manner which allows expression of the nucleic acid sequence.

To detect the phosphorylated product, any number of protocols known to those skilled in the art can be used including, but not limited to, Western blot analysis and apoptosis analysis. Antibodies, both monoclonal and polyclonal, can be made against epitopes derived from the site on the JNK substrate bound to a phosphate group. A SDS-PAGE procedure can be performed on homogenized cell extract and subsequently

subjected to Western blot analysis using an antibody specific for a phosphorylated JNK substrate, such as c-Jun.

An apoptosis analysis can also be performed in order to determine what effect, if any, the putative inhibitory agent has on JNK-associated activity. For example, an expression vector encoding JNK3 is transfected into an appropriate target cell to induce apoptosis. Target cells are cells that are susceptible to apoptosis. Rat hippocampal neuronal cell line HN33 is a preferred target cell. Alternatively, target cells which naturally contain JNK can be used. In either event, the target cells are cultured in the presence or absence of a test agent and the occurrence of apoptosis determined using known techniques. For example, staining the cell with Hoechst 3342 (Sigma Chemical Co.) and observing the stained cell under the microscope. Apoptotic cells appear containing clearly segmented, condensed chromatin. Alternatively, apoptosis can be determined by using the TUNEL assay as described by Thomas, L.B., *et al.*, *Exp. Neurol.*, 133:265-272 (1995). See also U.S. Patent No. 5,593,879, for techniques for examples of stains used to distinguish apoptotic cells.

In another embodiment, the invention pertains to a method for screening potential inhibitory agents of JNK activity, including JNK1, JNK2 and JNK3, by administering to an animal, including mammals, the agent and determining what effect, if any, the agent has on the animal's physiological status. The animal is given an amount of test agent sufficient to allow for proper pharmacodynamic absorption and tissue distribution in the animal. Preferably, the animal used is an example of a model system mimicking a neurological condition. However, to test the safety of the putative agent, a normal animal is preferably also subjected to the treatment. Following administration of the agent, the animal can be sacrificed and tissue sections from the brain, as well as other tissues, can be harvested and examined for apoptosis using, for example, the TUNEL assay. Yang, D.D., *et al.*, *Nature*, 389:865-870 (1997). In another embodiment, an animal model afflicted with a neurological condition (e.g., neurodegenerative disorder) can be administered a JNK and/or MLK inhibitor and the symptoms associated with the

neurological condition are evaluated. Attenuation, amelioration or improvement of the neurodegenerative symptoms can be assessed, whereby improvement is indicative of the inhibitors ability to prevent and/or treat the neurological condition.

The methods described above can be likewise employed to identify/screen for inhibitory agents of MLK-associated activity, including MLK1, MLK2 and MLK3. Appropriate MLK substrates include, but are not limited to, ATP and SEK1, a protein known to activate JNKs by phosphorylation. The MLK polypeptides useful in the methods of the present invention are preferably wild-type whose sequence is known and readily available. The human MLK2 polypeptide is described by Dorow, D.S., *et al.*, *Eur. J. Biochem.*, 234:491-500 (1995). Another MLK protein useful in the methods of the invention is described in GenBank Accession No. L32976.

The JNK and MLK useful in the methods of the invention are not limited to the naturally occurring sequences described above. JNK and MLK containing substitutions, deletions, or additions can also be used, provided that those polypeptides retain at least one activity associated with the naturally occurring polypeptide and are at least 70% identical to the naturally occurring sequence.

To determine the percent identity of two polypeptide sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid sequence). The amino acid residues at corresponding amino acid positions are then compared. When a position in the first sequence is occupied by the same amino acid residue as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions x 100).

The determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:5873-5877 (1993). Such an algorithm is

incorporated into the NBLAST and XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequence homologous to protein molecules useful in the methods of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, *Nucleic Acids Res*, 25:3389-3402 (1997).

- 5 When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers *et al.*, CABIOS (1989). Such an algorithm is incorporated into the ALIGN
- 10 program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

- The percent of identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In
- 15 calculating percent identity, only exact matches are counted.

- An example of a JNK or MLK that is not naturally occurring, though useful in the methods of the invention, is a JNK-gluthathione-S-transferase (JNK-GST) fusion protein. Such a protein can be produced in large quantities in bacteria and isolated. The JNK fusion protein can then be used in an *in vitro* kinase assay in the presence or
- 20 absence of a candidate drug for treating neurological conditions.

- The present invention also pertains to methods for testing the effectiveness of inhibitory compound identified using the methods of this invention for neuronal protection for the prevention and/or treatment of a variety of neurological disorders. In one embodiment of the invention, the effectiveness of neuronal protection by JNK or
- 25 MLK inhibitors against excitotoxicity stimuli can be assessed by the pre-treatment of HN33 cells with the identified JNK or MLK inhibitors prior to stimulation with glutamate, or other neurotoxins such as kainic acid, MPP, quinolinic acid or transfection of the mutated form of huntingtin or amyloid precursor protein (APP). For example,

HN33 cells, plated on a 96-well plate, are grown in DMEM-F12 medium and prior to experimentation, the medium is removed and cells are washed once with serum-free medium. 0.5 ml of serum free medium is added and cells are incubated at 37° C cell culture incubator for 10 minutes. In the case of the treatment with glutamate or other neurotoxins to induce apoptosis, the identified JNK or MLK inhibitor is added to the serum-free medium and incubation continued for another 5 minutes. Then glutamate or neurotoxin are added to the medium. If the JNK or MLK inhibitor is effective, the amount of apoptotic cells will be significantly reduced or totally inhibited, as compared with the appropriate control. Such a result indicates that these inhibitors are effective for the prevention of neuronal death in various neurological disorders. In the case of expression of mutated huntingtin or APP-C-100 (vector which expresses the c-terminal 100 amino acids of APP) to induce apoptosis, the JNK or MLK inhibitor is added during or 2-6 hours after transfection. In a particular embodiment, the IC<sub>50</sub> of the JNK or MLK inhibitors in suppression of neuronal apoptosis can be also assessed by this 96-well based assay. In this case, different concentrations of JNK or MLK inhibitor are added to the medium prior to the treatment with glutamate or other neurotoxins or during transfection of the mutated huntingtin or APP to establish a pharmacological profile for each inhibitor. The IC<sub>50</sub> of each inhibitor is a very important value for designing further study of the effectiveness in different animal models and for directing clinical trials of these inhibitors.

The present invention also pertains to methods for the prevention or treatment of neurological conditions, either through prophylactic administration prior to the occurrence of an event known to cause a neurological condition or therapeutic administration immediately following the event and periodically thereafter. Such prophylactic and therapeutic treatments are intended to prevent neuronal cell death or reduce the degree of cell death. Given the involvement of JNK and MLK in the cascade leading to neuronal cell death, these two kinases present targets for a therapeutic regime. According to the method, a mammal, including human, is administered an effective

therapeutic amount of an agent that targets JNK- and/or MLK-associated activity. A therapeutic amount for a given agent is that amount administered to achieve the desired result, for example, the inhibition of kinase activity in either JNK or MLK or both, or attenuation, amelioration of or improvement in the symptoms associated with the neurological condition.

In one embodiment, the JNK-associated activity that is targeted is JNK's kinase activity. By inhibiting JNK's activity with an agent, neuronal cell death can be avoided. The JNK activity to be targeted includes JNK1, JNK2 and JNK3. In another embodiment, the enzyme activity targeted is MLK. If MLK is not inhibited, then it will directly bind to and phosphorylate SEK1 resulting in its activation which in turn will stimulate JNK, thereby causing neuronal cell death. By inhibiting MLK activity, including MLK1, MLK2 and MLK3, the SEK1 phosphorylation and concomitant stimulation can be eliminated, thereby saving neuronal cells from apoptosis. This therapeutic approach can be used to prevent and/or treat neurological conditions, as described above. The inhibitory agents identified using the methods described herein are particularly useful for suppressing glutamate receptor-induced activation of JNK, glutamate-mediated toxicity and apoptosis caused by excitotoxicity.

Compounds identified using the methods described herein are designed to selectively inhibit the neuronal isoform of kinase which is involved in neuronal loss in neurodegenerative diseases. These kinase inhibitors will selectively decrease a specific kinase activity in neurons and protect neurons from a variety of oxidative stimuli thereby allowing a broad range of clinical applications. Because the neuronal isoform of kinase is selectively attenuated, side effects in peripheral tissues may be neglectable and because other isoforms of the kinase are present in neurons and will provide complementary function for the inhibited isoforms of the kinase, side effects in the central nervous system may also be minimal. A specific inhibitor of MLK2 or JNK3 should be an effective, low toxic neuroprotective drug for the treatment of a wide range of neurodegenerative disorders. In particular, two different kinases on the same

signaling pathway can be targeted. These different kinase inhibitors with similar clinical effects will allow to develop a clinical protocol to avoid drug tolerance and provide a life-long treatment.

Inhibitory agents of JNK, MLK or both, identified according to the methods of this invention, can be administered subcutaneously, intravenously, parenterally, intraperitoneally, intradermally, intramuscularly, topically, enteral (for example, orally), rectally, nasally, buccally, vaginally, by inhalation spray, by drug pump or via an implanted reservoir in dosage formulations containing conventional non-toxic, physiologically (or pharmaceutically) acceptable carriers or vehicles.

In a specific embodiment, it may be desirable to administer the agents of the invention locally to a localized area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, transdermal patches, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes or fibers.

In a specific embodiment when it is desirable to direct the drug to the central nervous system, techniques which can opportunistically open the blood brain barrier for a time adequate to deliver the drug there through can be used. For example, a composition of 5% mannitolose and water can be used. The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically (or prophylactically) effective amount of the agent, and a physiologically acceptable carrier or excipient. Such a carrier includes, but is not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The carrier and composition can be sterile. The formulation should suit the mode of administration.

Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions (for example, NaCl), alcohols, gum arabic, vegetable oils, benzyl alcohols, polyethylene glycols, gelatin, carbohydrates such as lactose, amylose or starch, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid esters,

hydroxymethylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired, mixed with auxiliary agents, for example, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds.

The compositions, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, polyvinyl pyrrolidone, sodium saccharine, cellulose, magnesium carbonate, etc.

The compositions can be formulated in accordance with the routine procedure as a pharmaceutical composition adapted for intravenous administration to human beings.

Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water, saline or dextrose/water. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

For topical application, there are employed as nonsprayable forms, viscous to semi-solid or solid forms comprising a carrier compatible with topical application and having a dynamic viscosity preferably greater than water. Suitable formulations include but are not limited to solutions, suspensions, emulsions, creams, ointments, powders,



enemas, lotions, sols, liniments, salves, aerosols, etc., which are, if desired, sterilized or mixed with auxiliary agents, for example, preservatives, stabilizers, wetting agents, buffers or salts for influencing osmotic pressure, etc. The drug may be incorporated into a cosmetic formulation. For topical application, also suitable are sprayable aerosol preparations wherein the active ingredient, preferably in combination with a solid or liquid inert carrier material, is packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant, e.g., pressurized air.

The amount of agents which will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. In addition, *in vitro* or *in vivo* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances.

The invention will now be further and specifically described by the following examples which are not intended to be limiting in any way. All publications cited herein are incorporated by reference in their entirety.

## EXAMPLES

20

### EXAMPLE 1: *Developing a Model for Neurodegeneration Based on Huntingtin Expression*

To develop a cell-based system in which apoptosis can be induced, polyglutamine-expanded huntingtin was introduced into cell lines.

25

To assess whether polyglutamine-expanded huntingtin causes toxicity, full-length huntingtin expression constructs containing 16, 48 or 89 CAG repeats were generated by assembly of a combination of RT-PCR products from normal and human HD

lymphoblast and plasmid cDNA clones IT16L and IT15B, which are described in HD Collaborative Research Group, *Cell* 72:971-983 (1993).

To construct the huntingtin expression vector with 16, 48, or 89 CAG repeats, the first third of the full-length construct was derived by ligation of IT16LL (bp 932-3018) with three different PCR products (bp 2401-3270, bp 637-1429 and 187-858). A 3027 bp cDNA fragment was removed from the resulting construct and ligated to corresponding sites in the cDNA clones IT15B (bp3024-10366). The CAG repeat size in the full length huntingtin construct pFL16HD was 16. PCR products were generated from the genomic DNA of an adult patient with 48 CAG repeats and a juvenile onset case with 89 CAG repeats. These PCR products replaced the corresponding region in pFL16HD to generate the pFL48HD and pFL89HD with 48 and 89 CAG repeats, respectively. Colony hybridization and PCR were used to identify the 48 and 89 CAG huntingtin clones and the positive clones were verified by DNA sequence analysis.

The resulting constructs pFL16HD, pFL48HD or pFL89HD were transiently transfected into 293 embryonic kidney cells, and expression of huntingtin was analyzed by immunoblotting using the anti-huntingtin monoclonal antibody 4C8. Trottier, Y., *et al.*, *Nature Genet.*, 10:104-110 (1995).

For transient expression of normal and polyglutamine-expanded huntingtin, 50% to 60% confluent HN33 or 293 cells were washed once with serum free medium prior to transfection. Transfection was performed by using lipofectin (Boehringer Mannheim) according to manufacturer instructions and fetal bovine serum was added to the medium 12 hours after transfection to a final concentration of 1%. Sixty µg of plasmid with 10µl of lipofectin /60 mm plate was used in all transfection experiments. After 72 hours, the transfection medium was removed and replaced by fresh medium with 1% fetal bovine serum. For immunoprecipitation and Western blotting, 293 cells were harvested 48-72 hours after transfection and lysed in 1% NP-40 lysate buffer. For the immunoprecipitation experiment, cell lysates were incubated with an affinity-purified anti-N-terminus huntingtin polyclonal antibody 437 for 4-6 hours. Liu, Y. F., *et al.*, *J.*

*Biol. Chem.*, 272:8121-8124 (1997). Cell lysates or precipitated proteins were resolved on SDS-PAGE, transferred and immunoblotted with an anti-huntingtin monoclonal antibody 4C8. All three huntingtin constructs constitutively expressed the huntingtin protein. Transfection of pcDNA1 (vector) or normal huntingtin with 16 CAG repeats (pFL16HD) did not lead to toxicity in HN33 cells. In addition, DNA fragmentation was not detectable using Tdt-mediated dUTP-biotin nick end labeling (TUNEL) assay. However, cell proliferation was slightly suppressed.

The TUNEL assay was performed by using a TUNEL assay kit (Boehringer Mannheim). HN33 cells were plated on a slide culture chamber. Transient transfection as conducted as described above. Transfection medium was removed at specified times post-transfection. The cells were washed once with serum free medium, fixed with 4% paraformaldehyde and then permeabilized with 0.1% of Triton X-100. The TUNEL assay was performed as described in the manufacturer's instructions provided with the kit.

Expression of mutated huntingtin with 48 or 89 CAG repeats (pFL48HD or pFL89HD) induced cell toxicity in HN33 cells. Apoptosis was initially observed between 20 and 24 hours after transfection by TUNEL staining. At 48 hours, about 75% of transfected cells were apoptotic.

A time-course of HN33 cell survival after transfection with pcDNA1, pFL16HD, pFL48HD or pFL89HD was performed forty eight hours after transfection, HN33 cells were fixed and stained with TUNEL. Living and apoptotic cells were counted in the high power field. The number of living cells in pcDNA1 transfectants is designated as 100%. Fig. 1 indicates that, cell toxicity induced by transfection of pFL89HD was slightly more severe than that mediated by pFL48HD, while transfection of pcDNA1 or pFL16HD did not produce measurable neuronal toxicity. Each data point in Fig. 1 is an average of four independent experiments.

To further confirm that the polyglutamine-expanded huntingtin-induced cell death described above was apoptotic, the interleukin 1 $\beta$  converting enzyme (ICE) inhibitor

zVAD-frm (Sarin *et al.*, *J. Exp. Med.* 184:2445-2449 (1996)) or CPP32 inhibitor zDEVD-frm (Rodriguez *et al.*, *J. Exp. Med.* 184:2067-2072 (1996)) was added to the medium during transfection to a inhibitor concentration of 10 µg/mL. Both inhibitors were obtained from Enzyme System Products, Inc. It was shown that ICE and CPP32 participated in apoptosis and that inhibitors zVAD-frm and zDEVD-frm blocked mutated huntingtin-mediated apoptotic cell death at 48 hours post-transfection (Fig. 2). ICE cleaves inactive CPP32 participated in apoptosis and that inhibitors zVAD-frm and zDEVD-frm blocked mutated huntington-mediated apoptotic cell death at 48 hours post-transfection (Fig. 2). (Vaux *et al.*, *Proc. Natl. Acad. Sci. USA*, 93:2239-2244 (1996)). It was known that ICE cleaves inactive CPP32 precursor thereby activating the enzyme. This result therefore suggests that expression of polyglutamine-expanded huntingtin stimulates ICE, which in turn activates CPP32 to induce apoptotic cell death.

#### 15 EXAMPLE 2: *Role of JNK in Neuronal Apoptosis*

Whether expression of polyglutamine-expanded huntingtin induces activation of JNK was investigated. GST c-Jun (1-89 aa) was utilized as a substrate to measure JNK activity in cell lysates from HN33 cells transfected with pcDNA1 (control), pFL16HD, pFL48HD or pFL89HD.

20 HN33 cells were lysed with 1% Triton buffer 16 hours after transfection. Cell lysates were incubated with glutathione-S-transferase (GST)-c-jun (1-89) fusion protein immobilized on glutathione sepharose beads to isolate JNK. These beads were resuspended in 30 µL kinase buffer. The kinase reaction was performed at 30°C for 30 minutes and then stopped by adding SDS sample buffer to the reaction. The reaction was analyzed by Western blotting using a phospho (ser63)-specific c-jun antibody (New England BioLabs).

A low level of JNK activation was observed in control cells. Transfection of a plasmid encoding normal huntingtin with 16 CAG repeats did not further increase the

amount of serine phosphorylated GST c-Jun and thus did not stimulate JNK activity. Expression of mutated huntingtin with 48 or 89 CAG repeats, however, significantly increased the levels of JNK activity. Serine phosphorylated GST c-Jun was increased 7- to 8-fold, similar to the level induced by 30 minutes of ultraviolet light irradiation.

5 These results indicated that polyglutamine repeat expansion in huntingtin activated JNK in HN33 cells (Fig. 3).

Whether activation of JNK is responsible for polyglutamine-expanded huntingtin-induced apoptotic cell death in HN33 cells was next examined. It was known that JNK is specifically activated by SEK1, which is a dual-specificity kinase that phosphorylates  
10 both tyrosine and threonine residues of JNK, thereby activating it Sanchez *et al.*, *Nature*, 380:75-79 (1994)). A dominant negative mutant of SEK1 (K54R), known to specifically block JNK activation (Lin *et al.*, *Science*, 268:286-290 (1995); Yan *et al.*, *Nature*, 372:798-800 (1994)) was used.

HN33 cells were transiently transfected with various expression plasmids,  
15 including the SEK1 and SEK1 (K54R) expression vectors described in Lin, A., *et al.*, *Science*, 268:286-290 (1995). Forty-eight hours after transfection, the cells were fixed and subjected to the TUNEL assay as described above. The number of living cells transfected with pcDNA1 was designated at 100%. The data is summarized in Fig. 4. Each data point represents an average of three independent experiments.

20 Transient expression of wild-type or dominant negative SEK1 alone had little effect on the proliferation and survival of HN33 cells. Co-expression of pcDNA1 with pEBG (SEK1 expression vector backbone control plasmid) also did not lead to cell toxicity. However, co-transfection of wild-type SEK1 vector with pFL48HD or pFL89HD did not affect neuronal toxicity induced by mutated huntingtin. Co-  
25 expression of dominant negative mutant SEK1 (K54R) significantly prevented apoptotic cell death induced by mutated huntingtin. At 48 hours after transfection, about 25% to 30% of cells had undergone apoptosis, compared to about 75% of cells containing pFL48HD or pFL89HD alone.

The foregoing data showed that a specific inhibitor of JNK, SEK1 (K54R), can inhibit polyglutamine-expanded huntingtin-induced apoptosis and may be a useful drug for treating a neurological condition. Thus it was possible to test a compound's potential as a drug by contacting the compound with a JNK inside a cell.

5 To determine whether the SEK1-JNK signal transduction pathway is involved in neuronal cell death induced by excitotoxicity induced by glutamate or kainate receptors, HN33 cells were transfected with pEBG (SEK1 expression vector backbone control plasmid), wild-type SEK1 expression vector, or dominant negative mutant SEK1 (K54R) expression vector. Forty-eight hours after transfection, the media covering the  
10 cells was replaced with fresh serum-free media supplemented without (control) or with 250  $\mu$ M glutamate or 200  $\mu$ M kainic acid (kainate). Cells were incubated in 37°C for six hours. Then cells were fixed and stained with TUNEL. TUNEL negative cells (living cells) were counted. The results in Fig. 5 show that SEK1 (K54R) blocks apoptosis induced by glutamate or kainate receptor activation. Each data point  
15 represents the average of three independent experiments.

To determine further the role of JNK in glutamate-induced neuronal apoptosis, HN33 cells were treated with glutamate (250  $\mu$ M) or kainic acid (200  $\mu$ M) at room temperature for 15 minutes and then lysed in 1% Triton X-100 buffer. Cell lysates were incubated with GST-c-Jun (1-89) fusion protein immobilized on glutathione sepharose  
20 beads to isolate JNK. These beads were resuspended in 30  $\mu$ l kinase buffer and the kinase reaction was performed at 30°C for 30 minutes and 10  $\mu$ l SDS sample buffer was added to stop the reaction. The samples were resolved by electrophoresis and transferred to a PVDF membrane (Millipore) and the JNK activity was analyzed by  
25 Western blotting using a phospho (Ser63)-specific c-Jun antibody (New England Biolabs). The results in Fig. 6 show that glutamate or kainate receptor activation induced elevation of the JNK activity. Each data point represents an average of three independent experiments.

To determine the role of activation of the SEK1-JNK signal transduction pathway in Alzheimer's diseases, a cDNA fragment encoding the full-length (Kang, J., *et al.*, *Nature*, 325:733-736 (1987)) or the last C-terminal 100 amino acid of amyloid precursor protein (APP) was inserted into pcDNA1 and the resulting plasmid was designated as APP-C-100. HN33 cells (60% of confluence) was transfected with pcDNA1 (control), wild-type APP, APP-C-100 by using lipofectin (Boehringer Mannheim) according to manufacture instruction. Thirty  $\mu$ l of lipofectin/50 mm plate was used in all transfection experiments. Twelve to seventy-two hours after transfection, HN33 cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% of Triton X-100. The TUNEL staining was performed by using a TUNEL staining kit (Boehringer Mannheim) as described in manufacture instructions provided with the kit. TUNEL negative cells were counted under light microscope. The results in Fig. 7 show that expression of APP-C-100 induced rapid neuronal apoptosis. Cell death was initially observed between twenty to twenty-four hours after transfection and at seventy-two hours, all cells were apoptotic, while expression of pcDNA1 (control) or wild-type APP did not induce neuronal cell death.

To determine the role of the JNK activity in neuronal cell death induced by APP-C-100, HN33 cells were transiently transfected with pcDNA1, wild-type APP or APP-C-100 using lipofectin as described above. After eighteen hours, cells were lysed in 1% Triton C-100 buffer. Cell lysates were incubated with GST-c-Jun (1-89) fusion protein immobilized on glutathione sepharose beads to isolate JNK. These beads were resuspended in 30  $\mu$ l kinase buffer and the kinase reaction was performed at 30°C for 30 minutes and 10  $\mu$ l SDS sample buffer was added to stop the reaction. The JNK activity was analyzed by Western blotting using a phospho (Ser63)-specific c-Jun antibody (New England BioLabs). The results in Fig. 8 show that expression of APP-C-100 but not wild-type APP stimulated the JNK activity in HN33 cells.

To further determine the role of activation of the JNK activation in Alzheimer's diseases, HN33 cells were transfected with pcDNA1, wild-type APP expression vector,

or mutant SEK1 expression vector using lipofectin (Boehringer Mannheim) according to manufacture instruction. Forty-eight hours after transfection, cells were fixed and stained with TUNEL. TUNEL negative cells (living cells) were counted. The results in Fig. 9 show that SEK1 (K54R) blocks apoptosis induced by expression of APP-C-100.

5 Taken together, these results indicate the elevation of the JNK activity is a common cause of neuronal cell death, regardless of the cause. Since excitotoxicity is a final common pathway for neuronal loss in neurodegenerative disease as well as in acute insults, inhibition of the JNK activity will prevent neuronal death in these neurological conditions.

### 10 EXAMPLE 3: *Role of MLK in Neuronal Apoptosis*

A kinase dead version of MLK2 was generated by introduction of A-G point mutation at position 651 (codon AGG to GAG) by overlapping extension using polymerase chain reaction with mutated oligonucleotides, to result in amino acid substitution of K to E in the ATP binding loop of the MLK2 kinase domain. Such a point mutation leads to total loss of kinase activity of MLK2 and a kinase dead version of MLK2 will act as a dominant mutant and inhibit MLK2 activation-mediated actions. Tibbles, L.A., *et al.*, *EMBO. J.*, 15:7026-7036 (1996). The cDNA fragment of wild-type or kinase dead MLK2 was inserted into pRK5CMV with a C-terminal myc tag.

20 Nagata, K., *et al.*, *EMBO. J.*, 17:149-158 (1998).

To examine whether expression of MLK2 induces neuronal cell death, HN33 cells were transfected with pRK5CMV, wild-type or kinase dead MLK2 expression vector using lipofectin (Boehringer Mannheim) according to manufacture instruction. Forty-eight hours after transfection, cells were fixed and stained with TUNEL, as described above. TUNEL negative cells (living cells) were counted. The results in Fig. 10 show that expression of MLK2 induced apoptosis in HN33 cells while expression of the kinase dead version of MLK2 did not generate any cell toxicity.



To determine whether the role of the SEK1-JNK signal transduction pathway in neuronal cell death induced by expression of MLK2, HN33 cells were co-transfected with pEBG + pRK5CMV (control), wild-type SEK1 expression vector + wild-type MLK2 expression vector, or dominant negative mutant SEK1 (K54R) expression vector + wild-type MLK2 expression vector. Forty-eight hours after transfection, cells were fixed and stained with TUNEL. TUNEL negative cells (living cells) were counted. The results in Fig. 11 show that SEK1 (K54R) attenuated apoptosis induced by MLK expression.

To determine the role of MLK2 in neuronal loss in Huntington's disease, HN33 cells were co-transfected with pEBG + pRK5CMV (control), normal huntingtin expression vector with 48 or 89 CAG repeats + kinase dead MLK2 expression vector. Forty-eight hours after transfection, cells were fixed and stained with TUNEL. TUNEL negative cells (living cells) were counted. The results in Fig. 12 show that co-expression of kinase dead MLK2 blocked neuronal apoptosis induced mutated huntingtin and demonstrated that the MLK-associated activity was involved in neuronal loss in Huntington's diseases.

The potential association of huntingtin with MLK2 was examined in 293 cells transfected with MLK2, normal huntingtin, or polyglutamine-expanded huntingtin in 293 cells at 48 hours after transfection. Expression of normal or mutated huntingtin failed to activate JNKs. Since 293 cells are rich in huntingtin (Liu, Y.F., *et al.*, *J Biol. Chem.*, 272:8121-8124 (1997)), the interaction of MLK2 with normal huntingtin in 293 cells was examined. C-Myc tagged MLK2 was transiently expressed in 293 cells, and MLK2 was precipitated with anti-c-myc tag 9E10 antibody (Santa Cruz).

C-myc-tagged MLK2 was transiently expressed in 293 cells, as described above, 48-72 hours after transfection, 293 cells were harvested and lysed in 1% NP-40 lysate buffer. Cell lysates were incubated with an anti-N-terminus huntingtin antibody 437 or 9E10 for 4-6 hours. The precipitated proteins were resolved on SDS-PAGE,

transferred, and immunoblotted with an anti-huntingtin monoclonal antibody 4C8, or anti-c-myc antibody 9E10.

In cell lysates from 293 cells transfected without or with pRK5CMV control vector described above, 9E10 failed to co-precipitate huntingtin. However, when the  
5 293 cells were transfected with c-myc tagged MLK2, huntingtin was easily detected in both 9E10 and 4C8 (anti-huntingtin) immunoprecipitates. Thus, MLK2 is associated with huntingtin.

Conversely, whether an anti-huntingtin antibody precipitated MLK2 was determined. Cell lysates from 293 cells with or without transfection of a MLK2  
10 expression vector were incubated with 437, an anti-huntingtin antibody, or 9E10. MLK2 was detectable in both 9E10 and 437 immunoprecipitates of 293 cell lysates transfected with MLK2. In non-transfected 293 cells, MLK2 was not found in either 9E10 or 437 immunoprecipitates. These results demonstrate that huntingtin is associated with MLK2 *in vivo*.

Next, we tried to examine the novel association of MLK2 with mutated huntingtin  
15 containing 48 CAG repeats in 293 cells. Expression of MLK2 or mutated huntingtin alone did not generate any cell toxic effect. Co-expression of MLK2 with normal huntingtin containing 16 CAG repeats also did not produce any cell toxicity while co-expression of mutated huntingtin with induced rapid cell death.

To determine whether the role of MLK2 in neuronal cell death induced by  
20 glutamate or kainate receptor activation, HN33 cells were transfected with pRK5CMV (control) or kinase dead MLK2 expression vector. Forty-eight hours after transfection, the media covering the cells was replaced with fresh serum-free media supplemented without (control) or with 250  $\mu$ M glutamate or 200  $\mu$ M kainate. Cells were incubated in  
25 37°C for six hours, fixed, and then stained with TUNEL. TUNEL negative cells (living cells) were counted. The results in Fig. 13 show that kinase dead MLK2 blocks apoptosis induced by glutamate or kainate receptor activation in HN33 cells.

To determine further the role of activation of the MLK activation in Alzheimer's diseases, HN33 cells were co-transfected with pcDNA1, wild-type expression vector, with kinase dead MLK2 expression vector. Forty-eight hours after transfection, cells were fixed and stained with TUNEL. TUNEL negative cells (living cells) were counted.

5 The results in Fig. 14 show that kinase dead MLK2 blocks apoptosis induced by expression of APP-C-100.

Taken together, these results shown that the MLK-associated activity mediated neuronal degeneration in Huntington's disease, Alzheimer's disease and excitotoxicity induced by glutamate or kainate receptor activation. Since excitotoxicity is a final  
10 common pathway for neuronal loss in neurodegenerative disease as well as in acute insults, inhibition of the MLK2 activity will prevent neuronal death in these neurological conditions.

#### EXAMPLE 4: *Preparation and purification of GST Fusion proteins*

15 Glutathione-S-transferase (GST)) c-Jun fusion protein was utilized as the substrate for JNK3 kinase assay and GST SEK1 fusion protein was utilized as the substrate for MLK2 kinase assay. To generate GST-c-Jun or GST-SEK1(K54R) fusion proteins, the cDNA fragment encoding the N-terminus of c-Jun 1-89 amino acid residues, or the full-length cDNA of SEK1, was subcloned into a pGEX vector. The  
20 vector was subsequently transformed into *E. coli* and the *E. coli* carrying pGEX-Jun or PGX-SEK1 was grown in LB medium. Expression of these GST fusion proteins was induced by 0.1 mM isopropyl-b-D-thiogalactopyraside (IPTG). Cells were pelleted and resuspended in 1/60 culture volume of MT PBS (150 mM NaCl, 16 mM Na<sub>2</sub>HPO<sub>4</sub>, 4 mM NaH<sub>2</sub>PO<sub>4</sub>, pH 7.3) and then lysed by mild sonication after adding Triton X-100 to  
25 the final concentration of 1% followed by centrifugation at 10,000 x g for 5 min at 4°C. The supernatant was mixed at room temperature in a 50 ml polypropylene tube on a rotating platform with 1-2 ml 50% glutathione agarose beads. After absorption for 2 min, beads were collected by brief centrifugation at 500 x g and washed three times with

50 ml MT PBS. The c-Jun or SEK1 GST fusion proteins were eluted by competition with free glutathione using 2 x 2 min washes twice with the same buffer, and stored. The purified GST fusion proteins were stored at -80°C in MT PBS at 4°C as a 50% solution.

5

EXAMPLE 5: *Transfection and JNK in vitro kinase assay*

For an *in vitro* JNK kinase assay, including JNK1, JNK2 and JNK3, isolated JNK protein was used. Isolation may be accomplished by chromatographic purification from tissue or molecular transfection of host cells followed by isolation. For molecular  
10 isolation of JNK, a full length cDNA of either JNK1, JNK2 or JNK3 glutathione-S-transferase (GST) fusion construct was inserted into pGEX-2T vector and expressed in a host, such as a bacterial cell (e.g., DH1 cell). The fusion protein was then purified from the host cell using standard techniques known to those skilled in the art. Isolation of the GST-fusion protein was accomplished using a glutathione affinity column.  
15 Approximately 5 µL of glutathione-Sepharose was used to recover the fusion protein. The resin was washed three times with lysis buffer (20 mM Tris-HCl, pH 8.0, 2 mM EDTA, 50 mM β-glycerophosphate, 1 mM Na<sub>2</sub>PO<sub>4</sub>, 1% Triton X-100, 10% glycerol, 1 mM phenylmethylsulfonyl fluoride (PMSF), 10 µg/mL leupeptin and 10 µg/mL aproptinin) and twice with detergent-free lysis buffer. The GST-fusion protein was  
20 eluted from the resin with 100 µL of 5 mM glutathione in a detergent-free lysis buffer.

The *in vitro* kinase reaction was carried out at 30°C for 15 minutes in a kinase mixture containing 0.5 µg of GST-JNK3, 1 µg of GST-c-Jun, 2.5 mM ATP and 5 µCi of [γ-<sup>32</sup>P] ATP with or without a test compound in 20 µL of kinase buffer (20 mM HEPES, pH 7.5, 15 mM MgCl<sub>2</sub>, 15 mM β-glycerophosphate, 0.1 mM Na<sub>2</sub>PO<sub>4</sub>, 2 mM  
25 dithiothreitol). The reaction was terminated by adding 20 µL of SDS sample buffer (Laemmli buffer). The phosphorylation was detected by SDS-PAGE, and the amount of <sup>32</sup>P incorporated was quantified with an image analyzer. Effectiveness of the test

compound as an inhibitor of JNK correlated with inhibitor of  $^{32}\text{P}$  incorporated GST-c-Jun.

EXAMPLE 6: *Transfection and MLK2 or MLK3 in vitro kinase assay*

- For an *in vitro* MLK 2 or 3 kinase assay, isolated MLK2 protein is used. Isolation
- 5 may be accomplished by chromatographic purification from tissue or molecular transfection of host cells followed by isolation. For molecular isolation of MLK, a full length cDNA of either MLK2 or MLK3 glutathione-S-transferase (hereinafter GST) fusion construct was inserted into pEGB vector and expressed in 293 host cells using lipofectin, as described above for transfection of mutated huntingtin into HN33 cells.
- 10 Forty eight hours after transfection, the cells were lysed using 0.5 ml at a lysis buffer (20 mM Tris-HCl, pH 8.0, 2mM EDTA, 50mM  $\beta$ -glycerophosphate, 1mM  $\text{Na}_2\text{VO}_4$ , 1% Triton X-100, 10% glycerol, 1mM phenylmethylsulfonyl fluoride (PMSF), 10  $\mu\text{g}/\text{ml}$  leupeptin and 10 $\mu\text{g}/\text{ml}$  aprotinin). The fusion protein was then purified from the host cell using standard techniques known to those skilled in the art. Isolation of the GST-
- 15 fusion protein was accomplished using a glutathione affinity column. Approximately 5  $\mu\text{L}$  of glutathione-Sepharose was used to recover the fusion protein. The resin was washed three times with lysis buffer and twice with detergent-free lysis buffer. The GST-fusion protein was eluted from the resin with 100  $\mu\text{L}$  of 5 mM glutathione in a detergent-free lysis buffer.
- 20 The *in vitro* kinase reaction was carried out at 30°C for 45 minutes in a 20  $\mu\text{L}$  kinase mixture containing 1  $\mu\text{g}$  of GST-MLK2 or MLK3, 1  $\mu\text{g}$  of GST-SEK1 (K54R), 2 mM ATP and 5  $\mu\text{Ci}$  or  $[\gamma\text{-}^{32}\text{P}]$  ATP in the absence or presence of a test compound in the kinase buffer (20 mM HEPES, pH 7.5, 15mM  $\text{MgCl}_2$ , 15mM  $\beta$ -glycerophosphate, 0.1 mM  $\text{Na}_2\text{VO}_4$ , 2mM dithiothreitol). The reaction was terminated by adding 20  $\mu\text{L}$  of
- 25 SDS sample buffer (Laemmli buffer). The phosphorylation was detected by SDS-PAGE, and the amount of  $^{32}\text{P}$  incorporated was quantified with an image analyzer. Effectiveness of the test compound as an inhibitor of MLK2 or MLK3 correlated with inhibition of  $^{32}\text{P}$  incorporated GST-SEK1 (K54R).

EXAMPLE 7: *MLK2 immunocomplex kinase assay*

An MLK2 immunocomplex kinase assay was used to screen for the MLK 2 or 3 inhibitor and to determine the specificity of the identified MLK inhibitor. The full-length cDNA of MLK 2 or 3 was inserted into pRK5 vector and tagged with c-myc at the C-terminus. 293 embryonic cells were grown in DMEM with 10% FBS in a 6-well dish and then were transfected with pRK5MLK2 using lipofectin as described above. Forty-eight hours after transfection, cells were lysed with 0.5 ml of a lysis buffer (20 mM Tris-HCl, pH 8.0, 2 mM EDTA, 50 mM  $\beta$ -glycerophosphate, 1 mM  $\text{Na}_2\text{VO}_4$ , 1% Triton X-100, 10% glycerol, 1mM phenylmethylsulfonyl fluoride (PMSF), 10  $\mu\text{g}/\text{ml}$  leupeptin, and 10 $\mu\text{g}/\text{ml}$  aproptinin) and the lysates were incubated with anti-c-myc tag antibody 9E10 for 1 hour and 20 $\mu\text{l}$  of a mixture (50 x 50) or protein G and A-Sepharose resin was add and continued to incubate for another 2-4 hours. The resin was washed twice with lysis buffer and twice with detergent-free lysis buffer. The pellet was then resuspended in 20  $\mu\text{l}$  of kinase buffer (20 mM HEPES, pH 7.5, 15 mM  $\text{MgCl}_2$ , 15mM  $\beta$ -glycerophosphate, 0.1 mM  $\text{Na}_2\text{VO}_4$ , 2 mM dithiothreitol, 25  $\mu\text{M}$  ATP, 5  $\mu\text{Ci}$  of [ $\gamma$ - $^{32}\text{P}$ ] ATP, and 0.5  $\mu\text{g}$  of SEK1(K54R) GST fusion proteins in the presence or absence (control) of 1 nM to 10  $\mu\text{M}$  of a mixture of compounds or the identified compounds. The kinase reaction was carried out at 30°C for 20 min. and stopped by adding 20  $\mu\text{l}$  of SDS sample buffer. The phosphorylation is detected by SDS-PAGE, and the amount of  $^{32}\text{P}$  incorporated is quantified with an image analyzer. If a compound is effective to inhibit MLKs, the amount of  $^{32}\text{P}$  incorporated is significantly reduced or totally inhibited.

EXAMPLE 8: *96 - Well cell based assay*

HN33 cells (~60% of confluence), plated on a 96-well plate, were grown in DMEM-F12 medium supplemented with 10% of fetal bovine serum (FBS). Prior to transfection, the medium was removed and cells were washed with serum-free medium once and 50  $\mu\text{l}$  of DMEM-F12 medium with 1% of FBS was added. The full-length

5 huntingtin expression plasmid containing 16 (control) ,48 or 89 CAG repeats or  
lipofectin solution were diluted with HBS first and mixed 1x1 volume and the mixture  
was incubated at room temperature for 15 min. Ten  $\mu$ l of the DNA-lipofectin mixture  
was added to the culture medium. A mixture of compound from a chemical library was  
added 6 hours after transfection. Twelve hours after transfection, additional FBS was  
added to the final concentration of 10%. Forty-eight hours after transfection, cells were  
washed with PBS once and fixed with 4% paraformaldehyde dissolved in PBS and  
permeabilized with 1% Triton X-100 in PBS containing 1% sodium citrate. Cell were  
then rinsed twice with PBS containing 1% BSA and cells were incubated with 25 $\mu$ l of  
10 TUNEL reaction solution for 1 hour at a 37°C cell culture incubator. Cell were rinsed  
with PBS for three times and 25 $\mu$ l of converter-AP solution was added and cells are  
returned to the incubator and incubated for 30 min. Cells were rinsed three times with  
PBS with 1% BSA and cells were incubated in a BCIP solution for 1-5 min at room  
temperature and rinsed three times with PBS and analyzed under light microscope.

15

While this invention has been particularly shown and described with references to  
preferred embodiments thereof, it will be understood by those skilled in the art that  
various changes in form and details may be made therein without departing from the  
spirit and scope of the invention as defined by the appended claims.

20

## CLAIMS

What is claimed is:

- 5      1.    A method for assessing a compound's ability to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, comprising:
- 10      a)    contacting a compound with neuronal cells having activated MLK and/or JNK activity;
- 10      b)    determining the number of neuronal cells that die;
- wherein a decreased number of dead neuronal cells in the presence of the compound compared to the number of dead neuronal cells in the absence of the compound is indicative of the compound's ability to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.
- 15      2.    The method of Claim 1, wherein the neuronal cells are transfected with a mutated protein or treated with a neurotoxin to induce apoptosis.
3.    The method of Claim 2, wherein the neuronal cells are HN33 cells.
- 20      4.    The method of Claim 2, wherein the mutated protein is polyglutamine stretch-expanded huntingtin or C-terminal 100 amino acids of amyloid precursor protein.
- 25      5.    The method of Claim 2, wherein the neurotoxin is glutamate, quinolinic acid or kainic acid.
6.    The method of Claim 1, wherein the neuronal cells are apoptotic neurons.



7. The method of Claim 1, wherein the neurological condition is a neurological disease whereby glutamate or kainic acid mediated excitotoxicity is involved in neuronal cell death.
- 5 8. The method of Claim 1, wherein the neurological condition is Huntington's disease or Alzheimer's disease.
9. A method for assessing a compound's ability to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition,  
10 comprising:
  - a) contacting a compound with neuronal cells transfected with a mutated protein or treated with a neurotoxin that induces neuronal cell death; and
  - b) determining the number of neuronal cells that die;wherein a decreased number of dead neuronal cells in the presence of the  
15 compound compared to the number of dead neuronal cells in the absence of the compound is indicative of the compound's ability to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.
10. The method of Claim 9, wherein the neuronal cells are HN33 cells.  
20
11. The method of Claim 9, wherein the mutated protein is polyglutamine stretch-expanded huntingtin or C-terminal 100 amino acids of amyloid precursor protein.
12. The method of Claim 9, wherein the neurological condition is a neurological  
25 disease whereby glutamate or kainic acid mediated excitotoxicity is involved in neuronal cell death.
13. The method of Claim 9, wherein the neurological condition is Huntington's disease or Alzheimer's disease.

14. A method for assessing the ability of a JNK and/or MLK inhibitor to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, comprising:
- a) contacting a JNK and/or MLK inhibitor with neuronal cells having activated MLK and/or JNK activity;
  - b) contacting, in the presence of the compound, surviving cells from step (a) with an agent that induces apoptosis; and
  - c) comparing the level of apoptosis in the cell in the presence of the JNK and/or MLK inhibitor with the level of apoptosis in the cell in the absence of the JNK and/or MLK inhibitor;
- wherein the JNK and/or MLK inhibitor is a potentially useful drug for treating the mammal when the level of apoptosis in the cell in the presence of the JNK and/or MLK inhibitor is less than the level of apoptosis in the cell in the absence of the JNK and/or MLK inhibitor.
15. The method of Claim 14, wherein the apoptotic agent is a neurotoxin.
16. The method of Claim 14, wherein the neurotoxin is glutamate, quinolinic acid or kainic acid.
17. The method of Claim 14, wherein step (b) is performed by transfecting the surviving neuronal cells with a mutated form of huntingtin or amyloid precursor protein.
18. The method of Claim 14, wherein the neuronal cells are HN33 cells.

19. A method for screening a compound's ability to inhibit JNK and/or MLK activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, comprising:
- a) contacting a compound with a JNK and/or MLK protein and substrate therefor;
  - b) measuring the level of JNK and/or MLK activity;
  - c) comparing the level of JNK and/or MLK activity in the presence of the compound with the level of JNK and/or MLK activity in the absence of the compound, wherein a decrease in JNK and/or MLK activity in the presence of the compound is indicative that the compound is a JNK and/or MLK inhibitor;
  - d) contacting the compound with neuronal cells having activated MLK and/or JNK activity;
  - e) comparing the occurrence of apoptosis in the cell in the presence of the compound with the occurrence of apoptosis in the cell in the absence of the JNK and/or MLK inhibitor;
- wherein the JNK and/or MLK inhibitor is a potentially useful drug for treating the mammal when the occurrence of apoptosis in the cell in the presence of the JNK and/or MLK inhibitor is less than the occurrence of apoptosis in the cell in the absence of the JNK and/or MLK inhibitor.
20. The method of Claim 19, wherein JNK is JNK1, JNK2 or JNK3 and MLK is MLK1, MLK2 or MLK3, or combinations thereof.
21. The method of Claim 20, wherein the JNK and/or MLK activity is a kinase activity.

22. The method of Claim 19, wherein the neurological condition is a neurological disease whereby glutamate or kainic acid mediated excitotoxicity is involved in neuronal cell death.
- 5 23. The method of Claim 19, wherein the neurological condition is Huntington's disease or Alzheimer's disease.
24. A method for assessing a compound's ability to inhibit JNK and/or MLK activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or  
10 having a neurological condition, comprising:  
a) incubating a compound in the presence of JNK and/or MLK and appropriate JNK and/or MLK substrate therefor, under conditions sufficient for enzymatic activity; and  
b) determining the presence or amount of phosphorylated product;  
15 wherein a change in amount of phosphorylated product, when compared to incubating JNK and/or MLK with appropriate substrates absent the compound, is indicative of the compound's ability to inhibit the enzymatic activity of JNK and/or MLK and thereby prevent neuronal cell death in a mammal susceptible to or having a neurological condition.
- 20 25. The method of Claim 24, wherein JNK is JNK1, JNK2 or JNK3 and MLK is MLK1, MLK2 or MLK3, or combinations thereof.
26. The method of Claim 24, wherein JNK substrates include c-Jun and a phosphate  
25 donor.
27. The method of Claim 24, wherein phosphorylated product of step (b) is phosphorylated c-Jun or phosphorylated SEK1.

28. The method of Claim 27, wherein the MLK substrates include SEK1 and a phosphate donor.
29. A method for assessing a compound's ability to inhibit JNK and/or MLK kinase activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, comprising:
- 5           (a)     contacting a neuronal cell with a compound under conditions sufficient for JNK and/or MLK enzymatic activity; and
- (b)     determining the presence or amount of phosphorylated JNK and/or MLK product;
- 10           wherein a change in amount of phosphorylated product, when compared to incubating a cell absent the compound, is indicative of the compound's ability to inhibit JNK and/or MLK kinase activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.
- 15           30. The method of Claim 29 further comprising:
- (c)     determining cell viability after step (a);
- wherein any increase in the cell's viability status relative to a control is indicative of the compound's ability to inhibit JNK and/or MLK kinase activity thereby
- 20           affecting the viability of the cell.
31. The method of Claim 29, wherein alteration in the cell's viability status includes apoptosis.
- 25           32. The method of Claim 29, wherein JNK is JNK1, JNK2 or JNK3 and MLK is MLK1, MLK2 or MLK3, or combinations thereof.

33. A method for assessing a compound's ability to inhibit MLK and/or JNK kinase activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, comprising:
- 5 (a) administering to an animal an amount of compound under conditions sufficient to allow for proper pharmacodynamic absorption and distribution thereof in the animal; and
- (b) determining the physiological status of the animal;
- 10 wherein a change in physiological status, when compared to an animal not administered the compound, is indicative of the compound's ability to inhibit MLK and/or JNK kinase activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.
34. The method of Claim 33, wherein MLK is MLK1, MLK2 or MLK3 and JNK is JNK1, JNK2 or JNK3, or combinations thereof.
- 15 35. The method of Claim 33, wherein the animal is a mammal.
36. A method for treating a neurological disorder in a mammal in need thereof, comprising administering to the mammal an effective therapeutic amount of a
- 20 compound that inhibits JNK and/or MLK activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.
37. The method of Claim 36, wherein the neurological condition is a neurological disease whereby glutamate or kainic acid mediated excitotoxicity is involved in neuronal cell death.
- 25 38. The method of Claim 36, wherein the neurological condition is Huntington's disease or Alzheimer's disease.

39. The method of Claim 36, wherein JNK is JNK1, JNK2 or JNK3 and MLK is MLK1, MLK2 or MLK3, or combinations thereof.
- 5 40. A method for preventing neuronal cell death in a mammal susceptible to or having a neurological condition, comprising administering to the mammal in need thereof an effective therapeutic amount of a compound that inhibits JNK and/or MLK activity in a neuronal cell and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.
- 10 41. The method of Claim 40, wherein the neurological condition is a neurological disease whereby glutamate or kainic acid mediated excitotoxicity is involved in neuronal cell death.
- 15 42. The method of Claim 40, wherein the neurological condition is Huntington's disease or Alzheimer's disease.
- 20 43. A method for treating a neurological disorder in a mammal in need thereof, comprising administering to the mammal an effective therapeutic amount of a compound that inhibits JNK and/or MLK activity and thereby prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, wherein the compound is identified by a method for assessing a compound's ability to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition, comprising:
- 25 a) contacting a compound with neuronal cells having activated MLK and/or JNK activity;
- b) determining the number of neuronal cells that die;
- wherein a decreased number of dead neuronal cells in the presence of the compound compared to the number of dead neuronal cells in the absence of the

compound is indicative of the compound's ability to prevent neuronal cell death occurring in a mammal susceptible to or having a neurological condition.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
222



METHOD FOR IDENTIFYING JNK AND MLK  
INHIBITORS FOR TREATMENT OF NEUROLOGICAL CONDITIONS

5

## ABSTRACT OF THE DISCLOSURE

10 The present invention describes methods for identifying compounds that inhibit JNK and MLK kinase activity as drugs for treating a mammal susceptible to or having a neurological condition. This invention also discloses methods for preventing neuronal cell death and treating neurological conditions that involve neuronal cell death, particularly neurodegenerative diseases characterized by glutamine or kainate mediated toxicity, such as Huntington's disease and Alzheimer's disease.

B E C O M E A P A T E N T




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14